

marine stingers

Catostylus

Distribution

The Australian species is found in Queensland, Victoria and NSW and into South East Asia.

Common names

Blubber

Size and appearance

Mushroom-shaped bell 5-30cm in diameter. They are a creamy white brown colour (blue if found further north). No tentacles but eight 'fronds' or 'frills' hang underneath. The sting causes minor skin irritation.

First aid


1. Remove casualty from water if safe to do so
2. DRSABCD
3. Wash area with sea water (not freshwater)
4. Keep casualty at rest and reassure
5. Do not allow rubbing of the sting area
6. Place casualty's stung area in hot water (no hotter than the rescuer can comfortably tolerate) for 20 minutes
7. If local pain is unrelieved by heat or if hot water is not available, apply a cold pack or ice in a bag
8. Send for medical aid if symptoms persist or covers a sensitive area (e.g the eyes) and seek assistance from lifesavers/lifeguards

Did you know?

- Blubbers are in a group of jellyfish called rhizostomes; other rhizostomes have been demonstrated to use a sun compass navigation to migrate
- Australia is home to many different species of blubbers — most are larger, colourful and give only minor stings
- Blubbers are often home to a lot of strange marine life including crabs, brittle stars, barnacles and fish and crustaceans




Catostylus mosaicus



Distribution in Australian waters



Catostylus sting



Size relative to human

For more information, visit marinestingers.com.au

SURF LIFE SAVING
QUEENSLAND

